

AfarTJ-22MyrTo5Myr.nb

Several points along the (mostly) non-extended edges of the Arabian, Nubian, and Somalian plates at the Afar triple junction have been selected and grouped with the rift that they are adjacent to: Red Sea rift, East African rift, Aden rift.

These points are moved back to where they were 22 million years ago, or about the time the East Africa rift started spreading.

We assume that the current angular velocity vectors for the Arabian, Nubian, and Somalian plates have been constant for the past 22 Myr and will continue to be constant in the next few million years, during the time we are interested in modeling.

Revised April 23, 2019

Input

Select the future time for the model

```
In[*]:= modelTimeF = 5;
```

Select the Euler pole data to be used for each of the three plates

```
In[*]:= poleLatSM = 54.6131;  
poleLongSM = -71.5876;  
angSpeedSM = 0.225471;
```

```
In[*]:= poleLatNB = 50.3991;  
poleLongNB = -36.5919;  
angSpeedNB = 0.189645;
```

```
In[*]:= poleLatAR = 42.155;  
poleLongAR = -0.270986;  
angSpeedAR = 0.456179;
```

Input the geographic data marking the edge of (mostly) undeformed continental crust for each of the three plates

The following matrices contain the geographic coordinates of several points along the (mostly) non-extended edges of the plate referenced in the matrix name. Each record corresponds to a different boundary point. The first column contains decimal latitudes ($^{\circ}$) and the second column contains decimal longitudes ($^{\circ}$). The data were collected by visual interpretation of a Mercator map projection of a global DEM presented by GeoMapApp (www.geomapapp.org).

```
In[ ]:= nubiaPtsToday =  $\begin{pmatrix} 12.662 & 39.507 \\ 11.613 & 39.590 \\ 10.367 & 39.814 \\ 9.674 & 39.744 \\ 8.69657 & 39.4531 \\ 7.0467 & 37.8161 \end{pmatrix}$ ;
```

```
In[ ]:= noRecsNB = 6;
```

```
In[ ]:= somaliaPtsToday =  $\begin{pmatrix} 7.2350 & 38.8180 \\ 8.8330 & 40.5044 \\ 9.440 & 41.493 \\ 9.518 & 42.047 \\ 9.748 & 43.128 \end{pmatrix}$ ;
```

```
In[ ]:= noRecsSM = 5;
```

```
In[ ]:= arabiaPtsToday =  $\begin{pmatrix} 17.095 & 43.124 \\ 15.441 & 43.370 \\ 14.179 & 43.576 \\ 13.693 & 45.035 \\ 13.885 & 45.756 \\ 14.218 & 46.986 \end{pmatrix}$ ;
```

```
In[ ]:= noRecsAR = 6;
```

User-Defined Functions

The function `unitVect3D` operates on a 3D vector (`vect_`) and outputs the Cartesian coordinates of the corresponding unit vector.

```
In[ ]:= unitVect3D[vect_] := {(vect[[1]]/Norm[vect]),  
  (vect[[2]]/Norm[vect]), (vect[[3]]/Norm[vect])};
```

```
In[ ]:= zRotation[w_, dT_] := {{Cos[(w dT) Degree], -Sin[(w dT) Degree], 0},  
  {Sin[(w dT) Degree], Cos[(w dT) Degree], 0}, {0, 0, 1}};
```

The function `geog2Cart` operates on two geographic coordinates (`lat_` and `long_`) and outputs the Cartesian coordinates of the unit location vector to that point.

```
In[ ]:= geog2Cart[lat_, long_] := {Cos[lat Degree] Cos[long Degree],  
  Cos[lat Degree] Sin[long Degree], Sin[lat Degree]};
```

The function `cart2Geog` operates on a 3D unit location vector (`vect_`) and outputs the geographic coordinates of that point in latitude and longitude.

```

In[ ]:= cart2Geog[vect_] :=
  Module[{lat, long, a, b, c, d, e, f}, a = ArcSin[vect[[3]]];
  b = {vect[[1]], vect[[2]], 0};
  c = If[ ((Abs[vect[[1]]] < (1 × 10-14)) && (Abs[vect[[2]]] < (1 × 10-14))),
    {1, 1, 0}, {vect[[1]]/Norm[b], vect[[2]]/Norm[b], 0}];
  d = {1, 0, 0};
  e = VectorAngle[c, d];
  f = If[(vect[[2]] < 0), (-e), (e)];
  lat = a (180/π);
  long = If[ ((Abs[vect[[1]]] < (1 × 10-14)) &&
    (Abs[vect[[2]]] < (1 × 10-14))), 0, (f (180/π))];
  {lat, long}];

```

The function `distAzFromVectors` operates on a 3D unit location vector to a reference point (`refPtVect_`) and a 3D unit location vector to another point (`otherPtVect_`), and outputs the distance in km and azimuth in degrees from the reference point to the other point.

```

In[ ]:= distAzFromVectors[refPtVect_, otherPtVect_] :=
  Module[{radEarth, circEarth, northPole, θ1, θ2, θ3, θ4, normVect1,
    normVect2, normVect3, distKm, bearing, answer}, radEarth = 6371.02;
  circEarth = 2 * π * radEarth;
  northPole = {0, 0, 1};
  normVect1 = Cross[northPole, refPtVect];
  normVect2 = Cross[otherPtVect, refPtVect];
  θ1 = VectorAngle[refPtVect, otherPtVect];
  distKm = (θ1 / (2 * π)) * circEarth;
  θ2 = (180/π) * (VectorAngle[normVect1, normVect2]);
  θ3 = (180/π) * (VectorAngle[normVect1, otherPtVect]);
  normVect3 = Cross[refPtVect, normVect1];
  θ4 = (180/π) * (VectorAngle[normVect3, otherPtVect]);
  bearing = If[ ((θ3 == 90) ∨ (θ3 == 180)),
    If[(θ4 <= 90), 0, 180], If[(θ3 > 90), (360 - θ2), θ2]];
  answer = {distKm, bearing};
  answer];

```

The function `circMotion` operates on a 3D unit location vector (`x_`), a coordinate transformation matrix (`m1_`), an angular speed (`w_`) and a time interval (`dT_`) in Myr, and outputs the unit location vector to that point after it rotates around its axis.

```

In[ ]:= circMotion[x_, m1_, w_, dT_] :=
  Module[{m2, m3, answer}, m2 = zRotation[w, dT];
  m3 = Inverse[m1];
  answer = m3.m2.m1.x;
  answer];

```

The function `spherePlot1` operates on a set of 3D unit location vectors of computed data (`inData_`) and two sets of 3D unit location vectors of colored markers (`markers1_` and

`markers2_`) to help establish the coordinate system(s), and outputs a graphic of a unit sphere on which the input points are plotted.

```

In[ ]:= spherePlot1[inData_, markers1_, markers2_] :=
  Module[{g1, g2, g3, g4, answer},
    g1 = Graphics3D[Sphere[{0, 0, 0}, 1], AspectRatio → 1,
      BoxRatios → {1, 1, 1}, PlotRange → All, PlotRangePadding → 0.1,
      ColorOutput → GrayLevel, Lighting → "Neutral"];
    g2 = ListPointPlot3D[inData, AspectRatio → 1, BoxRatios → {1, 1, 1},
      PlotStyle → Blue, PlotRange → All, PlotRangePadding → 0.1];
    g3 = ListPointPlot3D[markers1, AspectRatio → 1, BoxRatios → {1, 1, 1},
      PlotStyle → Red, PlotRange → All, PlotRangePadding → 0.1];
    g4 = ListPointPlot3D[markers2, AspectRatio → 1, BoxRatios → {1, 1, 1},
      PlotStyle → Green, PlotRange → All, PlotRangePadding → 0.1];
    answer = Show[g1, g2, g3, g4];
    answer];

In[ ]:= cycloid[obsΩ_, movΩ_, refPt_, t_] := Module[{a, b, c, d, e, f, g, h,
  i, j, k, l, m1, m2, m3, m4, m5, n, p, answer}, a = {1, 0, 0};
  b = {0, 1, 0};
  c = {0, 0, 1};
  d = unitVect3D[obsΩ];
  e = Norm[obsΩ];
  f = unitVect3D[movΩ];
  g = Norm[movΩ];
  h = unitVect3D[d × f];
  i = h;
  j = unitVect3D[h × d];
  k = unitVect3D[i × f];
  m1 = {{k[[1]], k[[2]], k[[3]]},
    {i[[1]], i[[2]], i[[3]]}, {f[[1]], f[[2]], f[[3]]}};
  m2 = {{Cos[(-g Degree) t], Sin[(-g Degree) t], 0},
    {-Sin[(-g Degree) t], Cos[(-g Degree) t], 0}, {0, 0, 1}};
  l = VectorAngle[d, k];
  If[l < (π/2), (n = 1), (n = (-1))];
  p = (n) (VectorAngle[f, d]);
  m3 = {{Cos[p], 0, -Sin[p]}, {0, 1, 0}, {Sin[p], 0, Cos[p]}};
  m4 = {{Cos[(e Degree) t], Sin[(e Degree) t], 0},
    {-Sin[(e Degree) t], Cos[(e Degree) t], 0}, {0, 0, 1}};
  m5 = {{j[[1]], h[[1]], d[[1]]}, {j[[2]], h[[2]], d[[2]]},
    {j[[3]], h[[3]], d[[3]]}};
  answer = m5.m4.m3.m2.m1.refPt;
  answer];

```

The user-defined function `northArrow` takes the average of all the unit location vectors of points along the plate boundary (`viewPoint`) and a user-defined scale (`nArrowScale`), generally ranging

from about 0.05 to 0.1, and finds a suitable endpoint for a vector pointing toward north in the output graphics.

```

In[ ]:= northArrow[viewPoint_, nArrowScale_] :=
  Module[{north, center, normal, northward, answer}, north = {0, 0, 1};
    center = {0, 0, 0};
    normal = Cross[north, viewPoint];
    northward = Cross[viewPoint, normal] * nArrowScale;
    answer = {center, northward};
    answer];

In[ ]:= midpoint[cartVect1_, cartVect2_] :=
  Module[{a, answer}, a = (cartVect1 + cartVect2) / 2;
    answer = {(a[[1]] / Norm[a]), (a[[2]] / Norm[a]), (a[[3]] / Norm[a])};
    answer];

In[ ]:= intersection[vect1_, vect2_, vect3_, vect4_] :=
  Module[{a, b, c, d, answer}, a = Cross[vect1, vect2];
    b = Cross[vect3, vect4];
    c = Cross[b, a];
    d = {(c[[1]] / Norm[c]), (c[[2]] / Norm[c]), (c[[3]] / Norm[c])};
    answer = If[(VectorAngle[d, vect2] > (π/2)), -d, d];
    answer];

In[ ]:= average3Vectors[vect1_, vect2_, vect3_] :=
  Module[{a, b, answer}, a = (vect1 + vect2 + vect3) / 3;
    answer = {(a[[1]] / Norm[a]), (a[[2]] / Norm[a]), (a[[3]] / Norm[a])};
    answer];

In[ ]:= addExplanation[plateName_, geogCoords_, modTime_] :=
  Module[{a, answer}, a = Table[
    Prepend[geogCoords[[i]], plateName], {i, 1, Length[geogCoords]}];
    answer = Table[Append[a[[i]], modTime], {i, 1, Length[a]}];
    answer];

```

Computation

The model time is 22 Myr ago (-22).

```
In[ ]:= modelTime = -22;
```

Move each set of points back to where they would have been at the selected time with a circular motion around the respective plate pole.

Define the axes of the geographic coordinate system

```
In[ ]:= xGeog = {1, 0, 0};
```

```
In[ ]:= yGeog = {0, 1, 0};
```

```
In[ ]:= zGeog = {0, 0, 1};
```

Nubian Plate

Convert the input pole location data for the pole to the Nubian plate to Cartesian coordinates

```
In[ ]:= poleEulerNB = geog2Cart[poleLatNB, poleLongNB];
```

Define the angular speed “ ω ” for the Nubian plate

```
In[ ]:=  $\omega$ NB = angSpeedNB;
```

Define the axes of the Euler pole coordinate system

```
In[ ]:= zPoleNB = poleEulerNB;
```

```
In[ ]:= xPoleNB = Cross[zPoleNB, zGeog];
```

```
In[ ]:= yPoleNB = Cross[zPoleNB, xPoleNB];
```

Find the Cartesian coordinates of the reference points on the Somalian plate

```
In[ ]:= cartNBtoday = Table[geog2Cart[nubiaPtsToday[[i, 1]],
  nubiaPtsToday[[i, 2]]], {i, 1, noRecsNB}];
```

Define the transformation matrix between the geographic coordinate system and the Euler pole coordinate system

```
In[ ]:= jNB = 
$$\begin{pmatrix} \text{Dot}[xGeog, xPoleNB] & \text{Dot}[yGeog, xPoleNB] & \text{Dot}[zGeog, xPoleNB] \\ \text{Dot}[xGeog, yPoleNB] & \text{Dot}[yGeog, yPoleNB] & \text{Dot}[zGeog, yPoleNB] \\ \text{Dot}[xGeog, zPoleNB] & \text{Dot}[yGeog, zPoleNB] & \text{Dot}[zGeog, zPoleNB] \end{pmatrix};$$

```

Rotate the reference point around the Euler pole by an angle equal to the angular speed multiplied by the time interval.

```
In[ ]:= rotPtsNB22mybp = Table[circMotion[
  cartNBtoday[[i]], jNB,  $\omega$ NB, modelTime], {i, 1, noRecsNB}];
```

```
In[ ]:= rotGeogNB22mybp = Table[
  cart2Geog[rotPtsNB22mybp[[i]]], {i, 1, Length[rotPtsNB22mybp]}];
```

```
In[ ]:= MatrixForm[%]
```

Out[]//MatrixForm=

```

$$\begin{pmatrix} 10.0987 & 36.4342 \\ 9.04881 & 36.5032 \\ 7.80018 & 36.7098 \\ 7.10808 & 36.6317 \\ 6.1343 & 36.3304 \\ 4.50588 & 34.6802 \end{pmatrix}$$

```

Somalian Plate

Convert the input pole location data for the pole to the Somalian plate to Cartesian

coordinates

```
In[ ]:= poleEulerSM = geog2Cart[poleLatSM, poleLongSM];
```

Define the angular speed “ ω ” for the Somalian plate

```
In[ ]:=  $\omega$ SM = angSpeedSM;
```

Define the axes of the Euler pole coordinate system

```
In[ ]:= zPoleSM = poleEulerSM;
```

```
In[ ]:= xPoleSM = Cross[zPoleSM, zGeog];
```

```
In[ ]:= yPoleSM = Cross[zPoleSM, xPoleSM];
```

Find the Cartesian coordinates of the reference points on the Somalian plate

```
In[ ]:= cartSMtoday = Table[geog2Cart[somaliaPtsToday[[i, 1]],
    somaliaPtsToday[[i, 2]]], {i, 1, noRecsSM}];
```

Define the transformation matrix between the geographic coordinate system and the Euler pole coordinate system

```
In[ ]:= jSM = 
$$\begin{pmatrix} \text{Dot}[x\text{Geog}, x\text{PoleSM}] & \text{Dot}[y\text{Geog}, x\text{PoleSM}] & \text{Dot}[z\text{Geog}, x\text{PoleSM}] \\ \text{Dot}[x\text{Geog}, y\text{PoleSM}] & \text{Dot}[y\text{Geog}, y\text{PoleSM}] & \text{Dot}[z\text{Geog}, y\text{PoleSM}] \\ \text{Dot}[x\text{Geog}, z\text{PoleSM}] & \text{Dot}[y\text{Geog}, z\text{PoleSM}] & \text{Dot}[z\text{Geog}, z\text{PoleSM}] \end{pmatrix};$$

```

Rotate the reference point around the Euler pole by an angle equal to the angular speed multiplied by the time interval.

```
In[ ]:= rotPtsSM22mybp = Table[circMotion[
    cartSMtoday[[i]], jSM,  $\omega$ SM, modelTime], {i, 1, noRecsSM}];
```

```
In[ ]:= rotGeogSM22mybp = Table[
    cart2Geog[rotPtsSM22mybp[[i]]], {i, 1, Length[rotPtsSM22mybp]}];
```

```
In[ ]:= MatrixForm[%]
```

Out[]//MatrixForm=

$$\begin{pmatrix} 4.50884 & 34.6802 \\ 6.13429 & 36.3305 \\ 6.75848 & 37.3018 \\ 6.8465 & 37.8505 \\ 7.09675 & 38.9196 \end{pmatrix}$$

Arabian Plate

Convert the input pole location data for the pole to the Arabian plate to Cartesian coordinates

```
In[ ]:= poleEulerAR = geog2Cart[poleLatAR, poleLongAR];
```

Define the angular speed “ ω ” for the Arabian plate

```
In[ ]:=  $\omega$ AR = angSpeedAR;
```

Define the axes of the Euler pole coordinate system

```
In[ ]:= zPoleAR = poleEulerAR;
```

```
In[ ]:= xPoleAR = Cross[zPoleAR, zGeog];
```

```
In[ ]:= yPoleAR = Cross[zPoleAR, xPoleAR];
```

Find the Cartesian coordinates of the reference points on the Arabian plate

```
In[ ]:= cartARtoday = Table[geog2Cart[arabiaPtsToday[[i, 1]],
    arabiaPtsToday[[i, 2]]], {i, 1, noRecsAR}];
```

Define the transformation matrix between the geographic coordinate system and the Euler pole coordinate system

```
In[ ]:= jAR = 
$$\begin{pmatrix} \text{Dot}[x\text{Geog}, x\text{PoleAR}] & \text{Dot}[y\text{Geog}, x\text{PoleAR}] & \text{Dot}[z\text{Geog}, x\text{PoleAR}] \\ \text{Dot}[x\text{Geog}, y\text{PoleAR}] & \text{Dot}[y\text{Geog}, y\text{PoleAR}] & \text{Dot}[z\text{Geog}, y\text{PoleAR}] \\ \text{Dot}[x\text{Geog}, z\text{PoleAR}] & \text{Dot}[y\text{Geog}, z\text{PoleAR}] & \text{Dot}[z\text{Geog}, z\text{PoleAR}] \end{pmatrix};$$

```

Rotate the reference point around the Euler pole by an angle equal to the angular speed multiplied by the time interval.

```
In[ ]:= rotPtsAR22mybp = Table[circMotion[
    cartARtoday[[i]], jAR, ωAR, modTime], {i, 1, noRecsAR}];
```

```
In[ ]:= rotGeogAR22mybp = Table[
    cart2Geog[rotPtsAR22mybp[[i]]], {i, 1, Length[rotPtsAR22mybp]}];
```

```
In[ ]:= MatrixForm[%]
```

Out[]//MatrixForm=

```

$$\begin{pmatrix} 12.2359 & 37.8551 \\ 10.5666 & 37.9244 \\ 9.29096 & 37.9974 \\ 8.66775 & 39.3755 \\ 8.79103 & 40.0994 \\ 9.00871 & 41.3338 \end{pmatrix}$$

```

Construct tables of the current and back-rotated points along the continental margins of the Nubian, Somalian, and Arabian Plates

```
In[ ]:= modTime = 0;
```

```
In[ ]:= plateName = "NubiaContEdge";
```

```
In[ ]:= nubiaContEdgeToday =
    addExplanation[plateName, nubiaPtsToday, modTime];
```

```
In[ ]:= plateName = "SomaliaContEdge";
```

```
In[ ]:= somaliaContEdgeToday =
    addExplanation[plateName, somaliaPtsToday, modTime];
```

```
In[ ]:= plateName = "ArabiaContEdge";
```

```
In[ ]:= arabiaContEdgeToday =
    addExplanation[plateName, arabiaPtsToday, modTime];
```

```
In[ ]:= Clear[plateName];
```



```

In[*]:= resultsContEdgeToday = Join[nubiaContEdgeToday,
    somaliaContEdgeToday, arabiaContEdgeToday];

In[*]:= modTime = -22;

In[*]:= plateName = "NubiaContEdge";

In[*]:= nubiaContEdge22mybp =
    addExplanation[plateName, rotGeogNB22mybp, modTime];

In[*]:= plateName = "SomaliaContEdge";

In[*]:= somaliaContEdge22mybp =
    addExplanation[plateName, rotGeogSM22mybp, modTime];

In[*]:= plateName = "ArabiaContEdge";

In[*]:= arabiaContEdge22mybp =
    addExplanation[plateName, rotGeogAR22mybp, modTime];

In[*]:= Clear[plateName];

In[*]:= resultsContEdge22mybp = Join[nubiaContEdge22mybp,
    somaliaContEdge22mybp, arabiaContEdge22mybp];

```

Rotate the current continental margins forward

```

In[*]:= rotPtsNB5Myr = Table[circMotion[
    cartNBtoday[[i]], jNB,  $\omega$ NB, modelTimeF], {i, 1, noRecsNB}];

In[*]:= rotGeogNB5Myr =
    Table[cart2Geog[rotPtsNB5Myr[[i]]], {i, 1, Length[rotPtsNB5Myr]}];

In[*]:= rotPtsSM5Myr = Table[circMotion[
    cartSMtoday[[i]], jSM,  $\omega$ SM, modelTimeF], {i, 1, noRecsSM}];

In[*]:= rotGeogSM5Myr =
    Table[cart2Geog[rotPtsSM5Myr[[i]]], {i, 1, Length[rotPtsSM5Myr]}];

In[*]:= rotPtsAR5Myr = Table[circMotion[
    cartARtoday[[i]], jAR,  $\omega$ AR, modelTimeF], {i, 1, noRecsAR}];

In[*]:= rotGeogAR5Myr =
    Table[cart2Geog[rotPtsAR5Myr[[i]]], {i, 1, Length[rotPtsAR5Myr]}];

```

Construct table of the forward-rotated points along the continental margins of the Nubian, Somalian, and Arabian Plates

```

In[*]:= plateName = "NubiaContEdge";

In[*]:= nubiaContEdge5Myr =
    addExplanation[plateName, rotGeogNB5Myr, modelTimeF];

In[*]:= plateName = "SomaliaContEdge";

```

```

ln[*]:= somaliaContEdge5Myr =
    addExplanation[plateName, rotGeogSM5Myr, modelTimeF];

ln[*]:= plateName = "ArabiaContEdge";

ln[*]:= arabiaContEdge5Myr =
    addExplanation[plateName, rotGeogAR5Myr, modelTimeF];

ln[*]:= Clear[plateName];

ln[*]:= resultsContEdge5Myr =
    Join[nubiaContEdge5Myr, somaliaContEdge5Myr, arabiaContEdge5Myr];

```

Finding the Initial Triple Junction

Red Sea Rift

```

ln[*]:= midpt1at22mybp = midpoint[rotPtsNB22mybp[[1]], rotPtsAR22mybp[[1]]];
ln[*]:= midpt2at22mybp = midpoint[rotPtsNB22mybp[[3]], rotPtsAR22mybp[[3]]];

```

East Africa Rift

```

ln[*]:= midpt3at22mybp = midpoint[rotPtsNB22mybp[[5]], rotPtsSM22mybp[[2]]];
ln[*]:= midpt4at22mybp = midpoint[rotPtsNB22mybp[[6]], rotPtsSM22mybp[[1]]];

```

Aden Rift

```

ln[*]:= midpt5at22mybp = midpoint[rotPtsSM22mybp[[5]], rotPtsAR22mybp[[6]]];
ln[*]:= midpt6at22mybp = midpoint[rotPtsSM22mybp[[3]], rotPtsAR22mybp[[4]]];

```

Intersections and mean triple junction

The method of finding the triple junction use here is to find the mean of the three riftline intersections. Consequently, no microplate is modeled to grow in this triple junction over time.

```

ln[*]:= tj1 = intersection[midpt4at22mybp,
    midpt3at22mybp, midpt2at22mybp, midpt1at22mybp];

ln[*]:= tj2 = intersection[midpt5at22mybp,
    midpt6at22mybp, midpt3at22mybp, midpt4at22mybp];

ln[*]:= tj3 = intersection[midpt1at22mybp,
    midpt2at22mybp, midpt6at22mybp, midpt5at22mybp];

ln[*]:= meanTJcart22mybp = average3Vectors[tj1, tj2, tj3];

```

```

In[*]:= riftTJpoints22mybp = {midpt1at22mybp, midpt2at22mybp,
    meanTJcart22mybp, midpt3at22mybp, midpt4at22mybp, midpt3at22mybp,
    meanTJcart22mybp, midpt6at22mybp, midpt5at22mybp};

In[*]:= nubiaPtsCart22mybp = {midpt1at22mybp, midpt2at22mybp,
    meanTJcart22mybp, midpt3at22mybp, midpt4at22mybp};

In[*]:= nubiaPtsGeog22mybp = Table[cart2Geog[nubiaPtsCart22mybp[[i]]],
    {i, 1, Length[nubiaPtsCart22mybp]};

In[*]:= somaliaPtsCart22mybp = {midpt4at22mybp, midpt3at22mybp,
    meanTJcart22mybp, midpt6at22mybp, midpt5at22mybp};

In[*]:= somaliaPtsGeog22mybp = Table[cart2Geog[somaliaPtsCart22mybp[[i]]],
    {i, 1, Length[somaliaPtsCart22mybp]};

In[*]:= arabiaPtsCart22mybp = {midpt1at22mybp, midpt2at22mybp,
    meanTJcart22mybp, midpt6at22mybp, midpt5at22mybp};

In[*]:= arabiaPtsGeog22mybp = Table[cart2Geog[arabiaPtsCart22mybp[[i]]],
    {i, 1, Length[arabiaPtsCart22mybp]};

```

Construct table of model points at 22 Myr BP

```

In[*]:= modTime = -22;

In[*]:= plateName = "nubia";

In[*]:= nubiaGeog22mybp =
    addExplanation[plateName, nubiaPtsGeog22mybp, modTime];

In[*]:= plateName = "somalia";

In[*]:= somaliaGeog22mybp =
    addExplanation[plateName, somaliaPtsGeog22mybp, modTime];

In[*]:= plateName = "arabia";

In[*]:= arabiaGeog22mybp =
    addExplanation[plateName, arabiaPtsGeog22mybp, modTime];

In[*]:= Clear[plateName];

In[*]:= resultsGeog22mybp =
    Join[nubiaGeog22mybp, somaliaGeog22mybp, arabiaGeog22mybp];

```

Prepare the graphic output

Parameters that affect the graphic output

Set the size of the north arrow on the graphic.

```
In[ ]:= northArrowScale = 0.1;
```

Choose the distance from Earth's center (expressed in Earth radii) from which the boundary will be viewed in the graphic. The value assigned to this variable must be greater than 1.

```
In[ ]:= viewDistance = 1.28;
```

```
In[ ]:= centerOfMap = geog2Cart[11.5, 42.0];
```

Variable `averageVv` is the point from which the boundary is viewed looking toward the center of Earth, as plotted using the built-in *Mathematica* function `Graphics3D`

```
In[ ]:= averageVv = viewDistance * centerOfMap;
```


Code that creates the initial graphic output

Note : Some useful (easily visible) built - in colors for use in the graphics include the following : Red, Green, Blue, Black, Gray, Cyan, Magenta, Brown, Orange, Pink, and Purple.

```
In[ ]:= boundaryOut22mybp =
```

```
Graphics3D[{Arrow[northArrow[averageVv, northArrowScale]], Black,
  Line[Table[rotPtsAR22mybp[[i]], {i, 1, Length[rotPtsAR22mybp]}]],
  Line[Table[rotPtsSM22mybp[[i]], {i, 1, Length[rotPtsSM22mybp]}]],
  Line[Table[rotPtsNB22mybp[[i]], {i, 1, Length[rotPtsNB22mybp]}]],
  Red, Line[Table[riftTJpoints22mybp[[i]],
    {i, 1, Length[riftTJpoints22mybp]}]], Blue, Dashed,
  Line[Table[cartARtoday[[i]], {i, 1, Length[cartARtoday]}]],
  Line[Table[cartSMtoday[[i]], {i, 1, Length[cartSMtoday]}]],
  Line[Table[cartNBtoday[[i]], {i, 1, Length[cartNBtoday]}]]],
  AspectRatio → Automatic, Boxed → False,
  ViewVector → {averageVv, {0, 0, 0}}];
```

```
In[ ]:= boundaryOut22mybp
```



The graphic shown above shows the Afar triple junction (red) at -22 MYBP.

- The original triple junction at 22 MYBP is shown by the solid red lines in the lower left, and the solid black lines on either side are the edges of the current non-extended continental crust as rotated back over 22 Myr.
- The blue dashed lines are the edges of the current non-extended continental crust today.

```
In[ ]:= ClearAll[tj1, tj2, tj3];
```

Growing the triple junction from 22 MYBP to today

```
In[ ]:= modelTime0 = 22;
```

Nubian Plate

The following table contains the latitude (first column) and longitude (second column) for the 5 model points on the Nubian plate 22 million years ago. The third record is the modeled location of the triple junction at that time.

```
In[ ]:= MatrixForm[nubiaPtsGeog22mybp]
```

```
Out[ ]//MatrixForm=
```

$$\begin{pmatrix} 11.1682 & 37.142 \\ 8.5461 & 37.3523 \\ 7.46535 & 37.58 \\ 6.13429 & 36.3305 \\ 4.50736 & 34.6802 \end{pmatrix}$$

Rotate the reference point around the Euler pole by an angle equal to the angular speed

multiplied by the time interval.

```
In[ ]:= rotPtsNB0 =
Table[circMotion[nubiaPtsCart22mybp[[i]], jNB, ωNB, modelTime0],
{i, 1, Length[nubiaPtsCart22mybp]}];
```

Somalian Plate

The following table contains the latitude (first column) and longitude (second column) for the 5 model points on the Somalian plate 22 million years ago. The third record is the modeled location of the triple junction at that time.

```
In[ ]:= MatrixForm[somaliaPtsGeog22mybp]
Out[ ]//MatrixForm=
```

$$\begin{pmatrix} 4.50736 & 34.6802 \\ 6.13429 & 36.3305 \\ 7.46535 & 37.58 \\ 7.71436 & 38.3363 \\ 8.05449 & 40.1239 \end{pmatrix}$$

Rotate the reference point around the Euler pole by an angle equal to the angular speed multiplied by the time interval.

```
In[ ]:= rotPtsSM0 =
Table[circMotion[somaliaPtsCart22mybp[[i]], jSM, ωSM, modelTime0],
{i, 1, Length[somaliaPtsCart22mybp]}];
```

Arabian Plate

The following table contains the latitude (first column) and longitude (second column) for the 5 model points on the Arabian plate 22 million years ago. The third record is the modeled location of the triple junction at that time.

```
In[ ]:= MatrixForm[arabiaPtsGeog22mybp]
Out[ ]//MatrixForm=
```

$$\begin{pmatrix} 11.1682 & 37.142 \\ 8.5461 & 37.3523 \\ 7.46535 & 37.58 \\ 7.71436 & 38.3363 \\ 8.05449 & 40.1239 \end{pmatrix}$$

Rotate the reference point around the Euler pole by an angle equal to the angular speed multiplied by the time interval.

```
In[ ]:= rotPtsAR0 =
Table[circMotion[arabiaPtsCart22mybp[[i]], jAR, ωAR, modelTime0],
{i, 1, Length[arabiaPtsCart22mybp]}];
```

Red Sea Rift

```
In[ ]:= midpt1Today = midpoint[rotPtsNB0[[1]], rotPtsAR0[[1]]];
```

```
In[ ]:= midpt2Today = midpoint[rotPtsNB0[[2]], rotPtsAR0[[2]]];
```

East Africa Rift

```
In[ ]:= midpt3Today = midpoint[rotPtsNB0[[4]], rotPtsSM0[[2]]];
```

```
In[ ]:= midpt4Today = midpoint[rotPtsNB0[[5]], rotPtsSM0[[1]]];
```

Aden Rift

```
In[ ]:= midpt5Today = midpoint[rotPtsSM0[[5]], rotPtsAR0[[5]]];
```

```
In[ ]:= midpt6Today = midpoint[rotPtsSM0[[4]], rotPtsAR0[[4]]];
```

Intersections and mean triple junction

The method of finding the triple junction use here is to find the mean of the three riftline intersections. Consequently, no microplate is modeled to grow in this triple junction over time.

```
In[ ]:= tj1 = intersection[midpt4Today, midpt3Today, midpt2Today, midpt1Today];
```

```
In[ ]:= tj2 = intersection[midpt5Today, midpt6Today, midpt3Today, midpt4Today];
```

```
In[ ]:= tj3 = intersection[midpt1Today, midpt2Today, midpt6Today, midpt5Today];
```

```
In[ ]:= meanTJcartToday = average3Vectors[tj1, tj2, tj3];
```

```
In[ ]:= riftTJpoints0 =
  {midpt1Today, midpt2Today, meanTJcartToday, midpt3Today, midpt4Today,
   midpt3Today, meanTJcartToday, midpt6Today, midpt5Today};
```

```
In[ ]:= nubiaPtsCartToday = {midpt1Today,
  midpt2Today, meanTJcartToday, midpt3Today, midpt4Today};
```

```
In[ ]:= nubiaPtsGeogToday = Table[cart2Geog[nubiaPtsCartToday[[i]]],
  {i, 1, Length[nubiaPtsCartToday]}];
```

The following table contains the latitude (first column) and longitude (second column) for the 5 model points on the Nubian plate today. The third record is the modeled location of the triple junction today.

```
In[ ]:= MatrixForm[nubiaPtsGeogToday]
```

Out[]//MatrixForm=

$$\begin{pmatrix} 14.8537 & 41.3544 \\ 12.2503 & 41.7206 \\ 11.0651 & 42.166 \\ 8.76515 & 39.9787 \\ 7.14112 & 38.3169 \end{pmatrix}$$

```
In[ ]:= somaliaPtsCartToday = {midpt4Today,
  midpt3Today, meanTJcartToday, midpt6Today, midpt5Today};
```

```
In[ ]:= somaliaPtsGeogToday = Table[cart2Geog[somaliaPtsCartToday[[i]]],
  {i, 1, Length[somaliaPtsCartToday]}];
```

The following table contains the latitude (first column) and longitude (second column)

for the 5 model points on the Somalian plate today. The third record is the modeled location of the triple junction today.

```
In[ ]:= MatrixForm[somaliaPtsGeogToday]
```

```
Out[ ]//MatrixForm=
```

$$\begin{pmatrix} 7.14112 & 38.3169 \\ 8.76515 & 39.9787 \\ 11.0651 & 42.166 \\ 11.511 & 43.3131 \\ 11.9189 & 45.1035 \end{pmatrix}$$

```
In[ ]:= arabiaPtsCartToday = {midpt1Today,
    midpt2Today, meanTJcartToday, midpt6Today, midpt5Today};
```

```
In[ ]:= arabiaPtsGeogToday = Table[cart2Geog[arabiaPtsCartToday[[i]]],
    {i, 1, Length[arabiaPtsCartToday]}];
```

The following table contains the latitude (first column) and longitude (second column) for the 5 model points on the Arabian plate today. The third record is the modeled location of the triple junction today.

```
In[ ]:= MatrixForm[arabiaPtsGeogToday]
```

```
Out[ ]//MatrixForm=
```

$$\begin{pmatrix} 14.8537 & 41.3544 \\ 12.2503 & 41.7206 \\ 11.0651 & 42.166 \\ 11.511 & 43.3131 \\ 11.9189 & 45.1035 \end{pmatrix}$$

Construct table of model points at their current locations

```
In[ ]:= modTime = 0;
```

```
In[ ]:= plateName = "nubia";
```

```
In[ ]:= nubiaGeogToday = addExplanation[plateName, nubiaPtsGeogToday, modTime];
```

```
In[ ]:= plateName = "somalia";
```

```
In[ ]:= somaliaGeogToday =
    addExplanation[plateName, somaliaPtsGeogToday, modTime];
```

```
In[ ]:= plateName = "arabia";
```

```
In[ ]:= arabiaGeogToday =
    addExplanation[plateName, arabiaPtsGeogToday, modTime];
```

```
In[ ]:= Clear[plateName];
```

```
In[ ]:= resultsGeogToday =
    Join[nubiaGeogToday, somaliaGeogToday, arabiaGeogToday];
```



Prepare the graphic output

Code that creates the graphic output

Note: Some useful (easily visible) built-in colors for use in the graphics include the following: Red, Green, Blue, Black, Gray, Cyan, Magenta, Brown, Orange, Pink, and Purple.

```
In[ ]:= boundaryOut22mybp =
Graphics3D[{Arrow[northArrow[averageVV, northArrowScale]],
  Black, Line[Table[rotPtsAR0[[i]], {i, 1, Length[rotPtsAR0]}]],
  Line[Table[rotPtsSM0[[i]], {i, 1, Length[rotPtsSM0]}]],
  Line[Table[rotPtsNB0[[i]], {i, 1, Length[rotPtsNB0]}]], Red,
  Line[Table[riftTJpoints0[[i]], {i, 1, Length[riftTJpoints0]}]],
  Blue, Dashed,
  Line[Table[cartARToday[[i]], {i, 1, Length[cartARToday]}]],
  Line[Table[cartSMtoday[[i]], {i, 1, Length[cartSMtoday]}]],
  Line[Table[cartNBtoday[[i]], {i, 1, Length[cartNBtoday]}]],
  Black, Dashed,
  Line[Table[rotPtsAR22mybp[[i]], {i, 1, Length[rotPtsAR22mybp]}]],
  Line[Table[rotPtsSM22mybp[[i]], {i, 1, Length[rotPtsSM22mybp]}]],
  Line[Table[rotPtsNB22mybp[[i]], {i, 1, Length[rotPtsNB22mybp]}]],
  Red, Dashed, Line[Table[riftTJpoints22mybp[[i]],
    {i, 1, Length[riftTJpoints22mybp]}]], AspectRatio → Automatic,
  Boxed → False, ViewVector → {averageVV, {0, 0, 0}}];
```

```
In[ ]:= boundaryOut22mybp
```



The graphic shown above shows the Afar triple junction (red) at -22 MYBP and today.

- The original triple junction at 22 MYBP is the dashed red line in the lower left, and the dashed black lines on either side are the edges of the current non-extended continental crust as rotated back over 22 Myr.
- The solid red lines in the middle of the diagram, with the black north arrow sticking out of them, is the current (model) triple junction. They are flanked by solid black lines, which are the displaced plate boundary from 22 MYBP (the red dashed lines). The blue dashed lines are the edges of the current non - extended continental crust today. The area between the solid black lines here is the new oceanic crust + extended continental crust of the last 22 Myr.

```
In[ ]:= ClearAll[tj1, tj2, tj3];
```

Growing the triple junction from Today to a Future Time

```
In[ ]:= modelTimeF
```

```
Out[ ]:= 5
```

Nubian Plate

Rotate the reference point around the Euler pole by an angle equal to the angular speed multiplied by the time interval.

```
In[*]:= rotPtsNBF =
  Table[circMotion[nubiaPtsCartToday[[i]], jNB,  $\omega$ NB, modelTimeF],
    {i, 1, Length[nubiaPtsCartToday]}];
```

Somalian Plate

Rotate the reference point around the Euler pole by an angle equal to the angular speed multiplied by the time interval.

```
In[*]:= rotPtsSMF =
  Table[circMotion[somaliaPtsCartToday[[i]], jSM,  $\omega$ SM, modelTimeF],
    {i, 1, Length[somaliaPtsCartToday]}];
```

Arabian Plate

Rotate the reference point around the Euler pole by an angle equal to the angular speed multiplied by the time interval.

```
In[*]:= rotPtsARF =
  Table[circMotion[arabiaPtsCartToday[[i]], jAR,  $\omega$ AR, modelTimeF],
    {i, 1, Length[arabiaPtsCartToday]}];
```

Red Sea Rift

```
In[*]:= midpt1F = midpoint[rotPtsNBF[[1]], rotPtsARF[[1]]];
In[*]:= midpt2F = midpoint[rotPtsNBF[[2]], rotPtsARF[[2]]];
```

East Africa Rift

```
In[*]:= midpt3F = midpoint[rotPtsNBF[[4]], rotPtsSMF[[2]]];
In[*]:= midpt4F = midpoint[rotPtsNBF[[5]], rotPtsSMF[[1]]];
```

Aden Rift

```
In[*]:= midpt5F = midpoint[rotPtsSMF[[5]], rotPtsARF[[5]]];
In[*]:= midpt6F = midpoint[rotPtsSMF[[4]], rotPtsARF[[4]]];
```

Intersections and mean triple junction

The method of finding the triple junction use here is to find the mean of the three riftline intersections. Consequently, no microplate is modeled to grow in this triple junction over time.

```
In[*]:= tj1 = intersection[midpt4F, midpt3F, midpt2F, midpt1F];
In[*]:= tj2 = intersection[midpt5F, midpt6F, midpt3F, midpt4F];
In[*]:= tj3 = intersection[midpt1F, midpt2F, midpt6F, midpt5F];
```

```
In[ ]:= meanTJcartF = average3Vectors[tj1, tj2, tj3];
```

```
In[ ]:= riftTJpointsF = {midpt1F, midpt2F, meanTJcartF,
  midpt3F, midpt4F, midpt3F, meanTJcartF, midpt6F, midpt5F};
```

```
In[ ]:= nubiaPtsCartF = {midpt1F, midpt2F, meanTJcartF, midpt3F, midpt4F};
```

```
In[ ]:= nubiaPtsGeogF =
  Table[cart2Geog[nubiaPtsCartF[[i]]], {i, 1, Length[nubiaPtsCartF]}];
```

The following table contains the latitude (first column) and longitude (second column) for the 5 model points on the Nubian plate after "modelTimeF" million years in the future. The third record is the modeled location of the triple junction at that time.

```
In[ ]:= MatrixForm[%]
```

```
Out[ ]//MatrixForm=
```

$$\begin{pmatrix} 15.7179 & 42.2955 \\ 13.1192 & 42.6962 \\ 11.8943 & 43.2192 \\ 9.3621 & 40.8122 \\ 7.7394 & 39.1466 \end{pmatrix}$$

```
In[ ]:= somaliaPtsCartF = {midpt4F, midpt3F, meanTJcartF, midpt6F, midpt5F};
```

```
In[ ]:= somaliaPtsGeogF = Table[
  cart2Geog[somaliaPtsCartF[[i]]], {i, 1, Length[somaliaPtsCartF]}];
```

The following table contains the latitude (first column) and longitude (second column) for the 5 model points on the Somalian plate after "modelTimeF" million years in the future. The third record is the modeled location of the triple junction at that time.

```
In[ ]:= MatrixForm[%]
```

```
Out[ ]//MatrixForm=
```

$$\begin{pmatrix} 7.7394 & 39.1466 \\ 9.3621 & 40.8122 \\ 11.8943 & 43.2192 \\ 12.3956 & 44.4373 \\ 12.8176 & 46.2301 \end{pmatrix}$$

```
In[ ]:= arabiaPtsCartF = {midpt1F, midpt2F, meanTJcartF, midpt6F, midpt5F};
```

```
In[ ]:= arabiaPtsGeogF = Table[
  cart2Geog[arabiaPtsCartF[[i]]], {i, 1, Length[arabiaPtsCartF]}];
```

The following table contains the latitude (first column) and longitude (second column) for the 5 model points on the Arabian plate after "modelTimeF" million years in the future. The third record is the modeled location of the triple junction at that time.

```
In[ ]:= MatrixForm[%]
```

```
Out[ ]//MatrixForm=
```

$$\begin{pmatrix} 15.7179 & 42.2955 \\ 13.1192 & 42.6962 \\ 11.8943 & 43.2192 \\ 12.3956 & 44.4373 \\ 12.8176 & 46.2301 \end{pmatrix}$$

Construct table of model points at a specified future time

```
In[*]:= modTime = 5;
In[*]:= plateName = "nubia";
In[*]:= nubiaGeogF = addExplanation[plateName, nubiaPtsGeogF, modTime];
In[*]:= plateName = "somalia";
In[*]:= somaliaGeogF = addExplanation[plateName, somaliaPtsGeogF, modTime];
In[*]:= plateName = "arabia";
In[*]:= arabiaGeogF = addExplanation[plateName, arabiaPtsGeogF, modTime];
In[*]:= Clear[plateName];
In[*]:= resultsGeogF = Join[nubiaGeogF, somaliaGeogF, arabiaGeogF];
```

Prepare the graphic output

Code that creates the graphic output

Note : Some useful (easily visible) built - in colors for use in the graphics include the following : Red, Green, Blue, Black, Gray, Cyan, Magenta, Brown, Orange, Pink, and Purple.

```

In[ ]:= boundaryOutF =
  Graphics3D[{Arrow[northArrow[averageVV, northArrowScale]],
    Black, Line[Table[rotPtsARF[[i]], {i, 1, Length[rotPtsARF]}]],
    Line[Table[rotPtsSMF[[i]], {i, 1, Length[rotPtsSMF]}]],
    Line[Table[rotPtsNBF[[i]], {i, 1, Length[rotPtsNBF]}]], Red,
    Line[Table[riftTJpointsF[[i]], {i, 1, Length[riftTJpointsF]}]],
    Blue, Dashed,
    Line[Table[cartARToday[[i]], {i, 1, Length[cartARToday]}]],
    Line[Table[cartSMtoday[[i]], {i, 1, Length[cartSMtoday]}]],
    Line[Table[cartNBtoday[[i]], {i, 1, Length[cartNBtoday]}]], Black,
    Dashed, Line[Table[rotPtsAR0[[i]], {i, 1, Length[rotPtsAR0]}]],
    Line[Table[rotPtsSM0[[i]], {i, 1, Length[rotPtsSM0]}]],
    Line[Table[rotPtsNB0[[i]], {i, 1, Length[rotPtsNB0]}]],
    Line[Table[rotPtsAR22mybp[[i]], {i, 1, Length[rotPtsAR22mybp]}]],
    Line[Table[rotPtsSM22mybp[[i]], {i, 1, Length[rotPtsSM22mybp]}]],
    Line[Table[rotPtsNB22mybp[[i]], {i, 1, Length[rotPtsNB22mybp]}]],
    Red, Dashed,
    Line[Table[riftTJpoints0[[i]], {i, 1, Length[riftTJpoints0]}]],
    Line[Table[riftTJpoints22mybp[[i]],
      {i, 1, Length[riftTJpoints22mybp]}]], AspectRatio -> Automatic,
    Boxed -> False, ViewVector -> {averageVV, {0, 0, 0}}];

```

```


In[ ]:= boundaryOutF

```

```

Out[ ]:=

```



The graphic shown above shows the Afar triple junction (red) at -22 MYBP, today, and 10 Myr in the future.

- The original triple junction at 22 MYBP is the dashed red line in the lower left, and the

dashed black lines on either side are the edges of the current non-extended continental crust as rotated back over 22 Myr.

- The red dashed lines in the middle of the diagram, with the black north arrow sticking out of them, is the current (model) triple junction. They are flanked by black dashed lines, which are the displaced plate boundary from 22 MYBP. The blue dashed lines are the edges of the current non - extended continental crust today. The area between the dashed black lines here is the new oceanic crust + extended continental crust of the last 22 Myr.

- The solid red lines to the upper right is the ridge - triple - junction system 10 Myr in the future. The solid black lines are the displaced ridge - triple - junction system of today as displaced on each of the three plates over 10 Myr of finite displacement.

```
In[ ]:= ClearAll[tj1, tj2, tj3];
```

Construct the consolidated table of model points

```
In[ ]:= outFile1 = Join[resultsGeog22mybp, resultsGeogToday, resultsGeogF,
  resultsContEdge22mybp, resultsContEdgeToday, resultsContEdge5Myr];
```

```
In[ ]:= MatrixForm[%]
```

```
Out[ ]//MatrixForm=
```

nubia	11.1682	37.142	-22
nubia	8.5461	37.3523	-22
nubia	7.46535	37.58	-22
nubia	6.13429	36.3305	-22
nubia	4.50736	34.6802	-22
somalia	4.50736	34.6802	-22
somalia	6.13429	36.3305	-22
somalia	7.46535	37.58	-22
somalia	7.71436	38.3363	-22
somalia	8.05449	40.1239	-22
arabia	11.1682	37.142	-22
arabia	8.5461	37.3523	-22
arabia	7.46535	37.58	-22
arabia	7.71436	38.3363	-22
arabia	8.05449	40.1239	-22
nubia	14.8537	41.3544	0
nubia	12.2503	41.7206	0
nubia	11.0651	42.166	0
nubia	8.76515	39.9787	0
nubia	7.14112	38.3169	0
somalia	7.14112	38.3169	0
somalia	8.76515	39.9787	0
somalia	11.0651	42.166	0
somalia	11.511	43.3131	0
somalia	11.9189	45.1035	0
arabia	14.8537	41.3544	0
arabia	12.2503	41.7206	0
arabia	11.0651	42.166	0
arabia	11.511	43.3131	0
arabia	11.9189	45.1035	0
nubia	15.7179	42.2955	5
nubia	13.1192	42.6962	5
nubia	11.8943	43.2192	5
nubia	9.3621	40.8122	5
nubia	7.7394	39.1466	5

	somalia	7.7394	39.1466	5
	somalia	9.3621	40.8122	5
	somalia	11.8943	43.2192	5
	somalia	12.3956	44.4373	5
	somalia	12.8176	46.2301	5
	arabia	15.7179	42.2955	5
	arabia	13.1192	42.6962	5
	arabia	11.8943	43.2192	5
	arabia	12.3956	44.4373	5
	arabia	12.8176	46.2301	5
	NubiaContEdge	10.0987	36.4342	-22
	NubiaContEdge	9.04881	36.5032	-22
	NubiaContEdge	7.80018	36.7098	-22
	NubiaContEdge	7.10808	36.6317	-22
	NubiaContEdge	6.1343	36.3304	-22
	NubiaContEdge	4.50588	34.6802	-22
	SomaliaContEdge	4.50884	34.6802	-22
	SomaliaContEdge	6.13429	36.3305	-22
	SomaliaContEdge	6.75848	37.3018	-22
	SomaliaContEdge	6.8465	37.8505	-22
	SomaliaContEdge	7.09675	38.9196	-22
	ArabiaContEdge	12.2359	37.8551	-22
	ArabiaContEdge	10.5666	37.9244	-22
	ArabiaContEdge	9.29096	37.9974	-22
	ArabiaContEdge	8.66775	39.3755	-22
	ArabiaContEdge	8.79103	40.0994	-22
	ArabiaContEdge	9.00871	41.3338	-22
	NubiaContEdge	12.662	39.507	0
	NubiaContEdge	11.613	39.59	0
	NubiaContEdge	10.367	39.814	0
	NubiaContEdge	9.674	39.744	0
	NubiaContEdge	8.69657	39.4531	0
	NubiaContEdge	7.0467	37.8161	0
	SomaliaContEdge	7.235	38.818	0
	SomaliaContEdge	8.833	40.5044	0
	SomaliaContEdge	9.44	41.493	0
	SomaliaContEdge	9.518	42.047	0
	SomaliaContEdge	9.748	43.128	0
	ArabiaContEdge	17.095	43.124	0
	ArabiaContEdge	15.441	43.37	0
	ArabiaContEdge	14.179	43.576	0
	ArabiaContEdge	13.693	45.035	0
	ArabiaContEdge	13.885	45.756	0
	ArabiaContEdge	14.218	46.986	0
	NubiaContEdge	13.2496	40.205	5
	NubiaContEdge	12.2008	40.2909	5
	NubiaContEdge	10.9554	40.5186	5
	NubiaContEdge	10.2622	40.4502	5
	NubiaContEdge	9.28405	40.1612	5
	NubiaContEdge	7.62988	38.5263	5
	SomaliaContEdge	7.84496	39.7679	5
	SomaliaContEdge	9.43583	41.4638	5
	SomaliaContEdge	10.0384	42.4569	5
	SomaliaContEdge	10.1139	43.0123	5
	SomaliaContEdge	10.3387	44.0963	5
	ArabiaContEdge	18.2689	44.2669	5
	ArabiaContEdge	16.6205	44.5528	5


```
 ArabiaContEdge 15.3632 44.7886 5  
 ArabiaContEdge 14.9077 46.2661 5  
 ArabiaContEdge 15.1144 46.9867 5  
 ArabiaContEdge 15.4721 48.2163 5
```

```
In[ ]:= Export[  
  "/Users/vince/Desktop/Ryley/PlotData4GeoMapApp.xlsx", outFile1]  
Out[ ]:= /Users/vince/Desktop/Ryley/PlotData4GeoMapApp.xlsx
```

